

DRUG	Alcohol	Marijuana	Nicotine	Stimulants	Hallucinogens	Depressants	Inhalants	Narcotics
1) Physical Dependence 2) Psychological Dependence 3) Tolerance	1) Moderate 2) Moderate 3) yes	1) Possible 2) Possible 3) Yes	1) High 2) Possible 3) Yes	1) High 2) High 3) Yes	1) Possible 2) No 3) Yes	1) High 2) High 3) Yes	1) Low/ Unknown 2) Possible 3) Yes	1) High 2) High 3) Yes
Possible Short-term Effects	Slurred speech, disorientation, drunken behavior	Euphoria, relaxed inhibitions, increased appetite, distorted time perception, disorientation	Increased excitation, euphoria, dilated pupils, increased pulse rate & blood pressure, insomnia, appetite loss	Increased wakefulness and physical activity and decreased appetite	Illusions & hallucinations, poor perception of time and distance	Slurred speech, disorientation, drunken-like behavior without odor of alcohol	Slurred speech, impaired coordination, euphoria, dizziness, possible hallucinations & delusions	Euphoria, drowsiness, respiratory depression, constricted pupils, nausea
Possible Long-Term Effects	Lowered resistance to disease/illness, increased risk of ulcers, heart disease & cancers of the liver, mouth, throat and stomach, increased risk of violence & injury	Respiratory problems, increased risk of obstructed airways, lung infections, and cancer of the respiratory tract and lungs	Chronic lung disease, heart disease, stroke, cancer of the lungs, larynx, esophagus, mouth, bladder, cervix, pancreas & kidneys	Violence, anxiety, confusion, insomnia. Possible paranoia, delusions, auditory hallucinations, and mood disturbances	Possible flashbacks, where aspects of the drug experience recur without taking the drug. Long-lasting psychosis possible.	Long-term use also can lead to increased tolerance and physical addiction.	Kidney abnormalities (glue or paint inhalation), liver damage from solvent or toluene use, memory & attention impairment, decreased non-verbal intelligence.	AIDS from needle use, reduced sex drive, drug impurities, limited vision, menstrual irregularity, chronic constipation, mood swings, breathing & heart problems, tremors, seizures, restlessness
Overdose Effects	Shallow respiration, clammy skin, dilated pupils, weak or rapid pulse, coma, possible death	Fatigue, paranoia, possible psychosis	Apathy, long periods of sleep, irritability, depression, disorientation	Can be lethal. Death can be caused by convulsions, circulatory & respiratory collapse & high body temperature	Longer, more intense "trip" episodes, psychosis, possible death	Shallow respirations, clammy skin, dilated pupils, weak and rapid pulse, coma and possible death	Death possible from heart failure, asphyxiation, or aspiration	Slow and shallow breathing, clammy skin, convulsions, coma, possible death

